

Abundance of substructure in clusters of galaxies

Priyamvada Natarajan¹ and Volker Springel²

ABSTRACT

It is widely recognized that cold dark matter models predict abundant dark matter substructure in halos of all sizes. Galaxy-galaxy lensing provides a unique opportunity to directly measure the presence and the mass of such substructures in clusters of galaxies. Here we present the mass function of substructures obtained from lensing in massive HST cluster lenses, and compare it with that obtained from high-resolution cosmological N-body simulations. We find excellent agreement in the slope and amplitude in the mass range $10^{11} - 10^{12.5} M_{\odot}$ probed by the observations, highlighting a significant success of the CDM theory. At lower mass scales, the simulations predict a large abundance of small substructures below our detection threshold with galaxy-galaxy lensing, with many of them being plausibly associated with faint cluster galaxies. Our results suggest that the CDM substructure abundance on the scale of clusters is in good agreement with present observational data.

Subject headings: cosmology: dark matter — gravitation: gravitational lensing

1. Introduction

Arguably the strongest challenges presently faced by the otherwise highly successful cold dark matter (CDM) paradigm for structure formation in the Universe occur on ‘small scales’, i.e. within individual dark matter halos. One of the most prominent of these problems is the long standing and controversially debated question whether rotation curves of low surface brightness galaxies can be accommodated in the CDM theory (e.g. McGaugh & de Blok 1998; Hayashi et al. 2003; Navarro et al. 2004). Another important issue is concerned with the amount of dark matter substructure theoretically predicted by CDM, and whether this is in conflict with the relative paucity of satellites observed in the Milky Way and the Local Group (e.g. Moore et al. 1999; Klypin et al. 1999; Stoehr et al. 2002).

¹Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520-8101, USA, priya@astro.yale.edu

²Max-Planck Institut für Astrophysik, Garching, Germany, volker@mpa-garching.mpg.de

Interestingly, both of these problems have only been recognized once N-body simulations reached high enough resolution for detailed studies of the phase-space structure of dark matter halos, allowing them to probe the inner dark matter cusp to sub-kpc scales, and to establish that dark halos are filled with a host of self-bound dark matter satellites, instead of being the smooth halos envisioned based on earlier low-resolution work. It has now become clear that the existence of substructure is a generic prediction of hierarchical structure formation in CDM models, where the assembly of collapsed mass proceeds via a merger hierarchy that allows some of the infalling dense lumps to survive as dynamically distinct substructure inside virialized halos until late times.

On the scales of ordinary galaxies like the Milky Way ($\sim 10^{12} M_{\odot}$), the number of bound satellites predicted by the simulations is much larger than the dozen or so satellites actually detected around our Galaxy. Comparing the observed velocity distribution function of these satellites with the predicted one (Moore et al. 1999; Klypin et al. 1999), this has been interpreted as evidence for a serious surfeit of dark matter satellites, and even considered to mark a “crisis” of the Λ CDM model. However, as Stoehr et al. (2002) have shown, there is significant uncertainty in the conversion of stellar line-of-sight velocity dispersion into peak circular velocities of dark matter substructures. Using high-resolution simulations of a Milky Way sized halo, Stoehr et al. in fact were able to show that all the known satellites of the Milky Way can be comfortably hosted by the most massive dark matter substructures expected for its halo. In this picture, there is however still a sea of small mass dark matter sub-halos which are largely devoid of stars. The leading hypothesis to explain this situation is to allude to baryonic processes of galaxy formation, inhibiting star formation preferentially in low mass halos. Proposed processes for such feedback include a photo-ionizing UV background or the expulsion of gas in shallow potential wells by supernova explosions (e.g. Bullock et al. 2000; Benson et al. 2002; Kravtsov et al. 2004).

If vast numbers of *dark* sub-halos exist, lensing might be the best way to detect them (e.g. Trentham et al. 2001). This is not only true for the scales of galaxies, but particularly for rich clusters of galaxies, where strong and weak lensing effects can be combined in a powerful way to construct detailed mass models for clusters. This allows in principle a direct mapping of the sub-halo mass function, thereby providing an important test of the CDM paradigm. Upon comparison with the observed galaxy luminosity function of clusters, it can also give valuable insights into the galaxy formation process.

In this letter, we present results of the first comparison between the lensing determined substructure mass function in clusters on mass scales ranging from about $10^{11} - 10^{12.5} M_{\odot}$ with that inferred from high resolution cosmological N-body simulations. To this end, we construct high resolution mass maps of galaxy clusters by applying galaxy-galaxy lensing

techniques where substructure inside clusters is detected and mapped using the anisotropies that they produce in the observed shear field. We compare our lensing measurements directly with results from high-resolution N-body simulations, allowing us to test the robustness of the CDM model and the associated hierarchical galaxy formation paradigm. Note that compared to the scales of galaxies, we expect many more dark matter structures to be visible optically in clusters, making the comparison of sub-halo mass functions less affected by uncertainties in the galaxy formation physics. Therefore, full consistency between the abundance of optically detected galaxies, substructure in CDM N-body models, and sub-halos detected by lensing can be asked for; establishing such a consistency can be viewed as a strong test of the theoretical paradigm. Below, we outline our methodology for mapping substructure in clusters. We then present the mass function derived from the maximum-likelihood analysis. This is followed by a comparison with the substructure detected in simulated clusters, and a discussion of our results. We adopt $h = 0.7$, $\Omega_0 = 0.3$ and $\Omega_\Lambda = 0.7$.

2. Quantifying substructure in clusters with gravitational lensing

We obtain the mass spectrum of clumps in a cluster by combining constraints from strong lensing observations (highly magnified multiply imaged systems) and weak lensing (the radially averaged tangential shear profile). To this end, we use a self-similar, parametric mass model to describe the cluster as a composite of a large-scale smooth mass distribution and several sub-clumps (Natarajan & Kneib 1997; Natarajan et al. 1998, 2002, 2004). These sub-halos are associated with bright, early-type galaxies in the cluster under the assumption that mass traces light. The local anisotropies in the shear field induced by the sub-halos in their vicinity is then used statistically to quantify the mass of the sub-clumps and their spatial extents. A likelihood method is used to retrieve characteristic halo parameters (e.g.

Cluster	z	σ_{0*} (km s ⁻¹)	r_{t*} (kpc)	M_{ap}/L_v (M _⊙ /L _⊙)	M* (10 ¹¹ M _⊙)	σ_{clus} (km s ⁻¹)	$\rho_{\text{clus}}(r = 0)$ (10 ⁶ M _⊙ kpc ⁻³)
A 2218	0.17	180 ± 10	40 ± 12	5.8 ± 1.5	~ 14	1070 ± 70	3.95
A 2390	0.23	200 ± 15	18 ± 5	4.2 ± 1.3	~ 6.4	1100 ± 80	16.95
AC 114	0.31	192 ± 35	17 ± 5	6.2 ± 1.4	~ 4.9	950 ± 50	9.12
Cl 2244–02	0.33	110 ± 7	55 ± 12	3.2 ± 1.2	~ 6.8	600 ± 80	3.52
Cl 0024+16	0.39	125 ± 7	45 ± 5	2.5 ± 1.2	~ 6.3	1000 ± 70	3.63
Cl 0054–27	0.58	230 ± 18	20 ± 7	5.2 ± 1.4	~ 9.4	1100 ± 100	15.84

Natarajan & Kneib 1997; Natarajan et al. 1998; Geiger & Schneider 1998). On applying these techniques to an ensemble of HST cluster lenses (results are presented in Table 1) we find that the spatial extents inferred are consistent with tidal stripping; early-type galaxies do possess dark halos that extend well beyond the light but these halos are more compact than those around field galaxies of equivalent luminosity.

In performing the likelihood analysis to obtain characteristic parameters for the sub-clumps in the cluster we assume that light traces mass. This is an assumption that is well supported by galaxy-galaxy lensing studies in the field (Wilson et al. 2001) as well as in clusters (Clowe & Schneider 2002; Hoekstra 2003). The individual galaxies and the smooth cluster component are modeled self-similarly with truncated pseudo-isothermal mass distributions (PIEMD). The parameters that characterize a truncated PIEMD are: a truncation radius r_t identified with the tidal radius, a core radius r_0 and a central velocity dispersion σ_0 . For the smooth component the values of these parameters are set by the observed strong lensing features, and for the galaxies, combined constraints from the strong lensing and the weak shear field determine the best-fit parameters for a fiducial galaxy halo. These values recovered from the maximum likelihood analysis are shown in Table 1. In order to relate the observed light distribution in the early-type cluster galaxies to their masses a set of physically motivated scaling laws are assumed:

$$\sigma_0 = \sigma_{0*} \left(\frac{L}{L_*} \right)^{1/4} ; \quad r_0 = r_{0*} \left(\frac{L}{L_*} \right)^\alpha ; \quad r_t = r_{t*} \left(\frac{L}{L_*} \right)^\alpha . \quad (1)$$

The total mass of the sub-halo associated with a galaxy of luminosity L is:

$$M \propto \sigma_{0*}^2 r_{t*} \left(\frac{L}{L_*} \right)^{1/2+\alpha} ; \quad \frac{M}{L} \propto \sigma_{0*}^2 r_{t*} \left(\frac{L}{L_*} \right)^{1/2-\alpha} . \quad (2)$$

Note however that for our choice of mass model and value adopted for the exponent α , the mass to light ratio is not constant with radius within an individual galaxy halo. The derived mass spectrum of sub-halos is not a strong function of α , as discussed in (Natarajan et al. 2004).

Dark halos are associated with the locations of bright, early-type cluster galaxies and the fiducial parameters for a typical halo are then extracted from the likelihood analysis. A high resolution mass model for the entire cluster is built using the strong lensing regime to constrain the inner region ($r \sim r_{\text{Einstein}}$), and the local anisotropies in the shear field are used to obtain properties of the galaxy halos around early-type cluster members. Since the procedure involves a scaled, self-similar mass model that is parametric, we obtain a mass estimate for the dark halos (sub-clumps) of the cluster galaxies as a function of their luminosity. This provides us with a clump mass spectrum. Note that tidal truncation

by the cluster causes these halo masses to be lower than that of equivalent field galaxies at comparable redshifts obtained from galaxy-galaxy lensing. The fraction of mass in the clumps is only 10-20% of the total mass of the cluster within the inner $500 h^{-1} \text{ kpc}$ of these high central density clusters. We are limited to this spatial scale as the lensing analysis was performed on HST-WFPC2 data with pointings at cluster centers. The remaining 80-90% of the cluster mass is consistent with being smoothly distributed (in lumps with mass $M < 10^{10} M_{\odot}$).

In Fig. 1, we show the mass function retrieved from galaxy-galaxy lensing for each of the five HST clusters. There is a low-mass cut-off in the observed clump spectrum, at around $10^{11} M_{\odot}$, which is due to observational limitations. The mass resolution of this technique is limited by the depth and field of view of the Wide Field Planetary Camera (WFPC2) aboard the Hubble Space Telescope, by the number of background galaxies per foreground lens, and the reliability with which shapes can be measured for the faintest background galaxies in the HST image of the central region. Unfortunately, this limits the number of reliably determined lumps per rich cluster to about 40, implying that only the massive end of the clump spectrum corresponding to the brightest cluster galaxies can be probed. However, despite the low number statistics, the individual cluster measurements show a marked rise in substructure abundance towards lower mass scales. This becomes particularly apparent once the clusters are stacked, as we have done in the lower right panel of Fig. 1.

3. Cluster substructure in high resolution Λ CDM simulations

We analyze substructure in high-resolution dark matter simulations of clusters formed in the Λ CDM model. Our set of simulations is taken from the study carried out by Springel et al. (2001a) of a single rich cluster of mass $8.4 \times 10^{14} h^{-1} M_{\odot}$, simulated in 4 steps of ever increasing resolution using the parallel tree-code GADGET (Springel et al. 2001b). In these simulations (referred to as ‘S1’ to ‘S4’), the particle mass resolution increases from 6.87×10^9 to $4.68 \times 10^7 h^{-1} M_{\odot}$, corresponding to 1.3×10^5 particles up to about 20 million within the virial radius, making the highest resolution simulation in this series one of the best resolved simulations of a single rich cluster carried out to date. This simulated cluster has a comparable central density and mass within the inner $500 h^{-1} \text{ kpc}$ as the massive HST cluster-lenses studied here.

Self-bound gravitational substructure is found with the algorithm SUBFIND (Springel et al. 2001a). It starts by determining locally over-dense dark matter substructure candidates in a fully adaptive fashion, and then subjects each of them to a gravitational unbinding procedure, such that a catalog of self-bound dark matter substructures results. In the lowest

resolution simulation S1 of the series, 118 substructures can be detected, a number that increases to more than 4600 in the high-resolution simulation S4. Note however that these additionally resolved small substructures are all of ever lower mass; already the low resolution simulation captures the correct number of massive satellites, with (at least on average) the correct mass.

The measured differential mass function of substructures appears to be a power law, $dN/dm \propto m^{-\alpha}$, with slope close to $\alpha \simeq -1.8$ (Springel et al. 2001a; Helmi et al. 2002; De Lucia et al. 2004). Interestingly, this is very close to the result by Lee (2004) who has attempted an analytic calculation of the sub-halo mass function based on modeling the complex dynamical history of galaxies in the cluster with a parameterized model to account for the effects of global tidal truncation (see also Taylor & Babul 2001). Given that we have also demonstrated that the spatial extents of substructures inferred from galaxy-galaxy lensing are consistent with the tidal stripping hypothesis, this lends support to the validity of a simple tidal-limit approximation. Similar to the lensing result, both the analytic model of Lee (2004) and the numerical simulations find that only about 10% of the sub-halo mass is bound in substructures, most of it in a handful of most massive sub-halos which dominate the cumulative mass in the substructures.

We note that numerical studies by De Lucia et al. (2004) find that the substructure mass function depends only weakly on the properties of the parent halo mass. Also, the mass fraction in substructure is relatively insensitive to the tilt and overall normalization of the primordial power spectrum (Zentner & Bullock 2003). Only for radically altered CDM models, for example by truncating small-scale power, Zentner & Bullock find that their models yield projected substructure mass fractions that are lower than the estimates from strong lensing.

4. Comparison

We now compare the galaxy-galaxy lensing results with the substructure mass function obtained from the N-body simulations. In Fig. 2, we show histograms for the distribution of substructure masses in the four simulations of Springel et al. (2001a) and contrast them individually with the stacked result for the clusters A2218, A2390, and Cl0054. Since the number of substructures in the observable mass range is quite small, we expect large system-to-system variations between different clusters. The stacking of clusters of similar mass that we applied here reduces the associated scatter somewhat.

Comparing the lensing result with the four simulated clusters (which also show some

numerical scatter amongst each other), we find broad general agreement in the mass range $10^{11} - 10^{12.5} M_{\odot}$, both in the amplitude and the slope of the substructure mass function. Given that no free parameter or scaling has been applied to obtain this match, the agreement is in fact remarkable.

However unlike the observations, the simulations show no low-mass cut-off. If this cut-off is entirely due to resolution effects, which appears plausible, the interpretation is that the observations only see the ‘tip of the iceberg’ of the substructure distribution as far as their number is concerned, even though they detect most of the mass in substructures. Fig. 2 gives also a hint that the simulations may systematically over-predict the mass of the most massive satellites, but we caution that such an effect could also be caused a systematic effect in the mass estimate based on the lensing technique, for example if the massive substructures preferentially correspond to halos that have fallen in most recently.

5. Discussion

On comparing the clump mass spectrum obtained for galaxy clusters from high-resolution N-body simulations of Λ CDM models to those obtained from galaxy-galaxy lensing in HST cluster lenses, we find excellent agreement. Despite the fact that the lensing analysis assumes that mass traces light and is only sensitive to a restricted mass range, it is clear that there is no substructure problem in CDM on mass scales spanning $10^{11} - 10^{12.5} M_{\odot}$. This is in sharp contrast to the situation on galactic scales, where the paucity of observed satellites when compared with the abundant dark matter substructure predicted by simulations has been characterized as a crises for CDM. While the severity of this problem has probably been overstated initially – in fact it may have partially gone away by now (StoeHR et al. 2002) – it is clear that CDM predicts a rich spectrum of dark matter substructure extending to very small masses, both for clusters and galactic halos. In clusters, up to several hundred of the most massive substructures can be directly identified with the luminous cluster galaxies, and semi-analytic models of galaxy formation show that this association leads to highly successful models for the population of cluster galaxies (Springel et al. 2001a). On the other hand, on galactic scales, a much larger number of truly dark satellites must be present according to the N-body models. Gravitational lensing is probably our best bet to detect such structures of small mass, and in fact, on these small scales, the observed flux anomalies in multiply-imaged quasar systems (Mao & Schneider 1998; Chiba 2002; Dalal & Kochanek 2002; Metcalf & Madau 2001; Metcalf & Zhao 2002; Mao et al. 2004) have been interpreted as evidence for significant substructure in the mass range $10^4 < M < 10^8 M_{\odot}$.

In this letter, we have shown that the clump mass function independently determined

from lensing (a technique that is unaffected by the dynamical state of the cluster) is in excellent agreement with that obtained in high resolution cosmological N-body simulations of clusters of galaxies in the Λ CDM model.

PN thanks her collaborators on the HST cluster-lenses project: Jean-Paul Kneib, Ian Smail and Richard Ellis.

REFERENCES

- Benson, A. J., Frenk, C. S., Lacey, C. G., Baugh, C. M., & Cole, S. 2002, MNRAS, 333, 177
- Bullock, J. S., Kravtsov, A. V., & Weinberg, D. H. 2000, ApJ, 539, 517
- Chiba, M. 2002, ApJ, 565, 17
- Clowe, D. & Schneider, P. 2002, A&A, 395, 385
- Dalal, N. & Kochanek, C. S. 2002, ApJ, 572, 25
- De Lucia, G., Kauffmann, G., Springel, V., White, S. D. M., Lanzoni, B., Stoehr, F., Tormen, G., & Yoshida, N. 2004, MNRAS, 348, 333
- Geiger, B. & Schneider, P. 1998, MNRAS, 295, 497
- Hayashi, E., Navarro, J. F., Power, C., Jenkins, A. R., Frenk, C. S., White, S. D. M., Springel, V., Stadel, J., & Quinn, T. R. 2003, submitted, astro-ph/0310576
- Helmi, A., White, S. D., & Springel, V. 2002, Phys. Rev. D, 66, 063502
- Hoekstra, H. 2003, MNRAS, 339, 1155
- Klypin, A., Kravtsov, A. V., Valenzuela, O., & Prada, F. 1999, ApJ, 522, 82
- Kravtsov, A. V., Gnedin, O. Y., & Klypin, A. A. 2004, ApJ, 609, 482
- Lee, J. 2004, ApJ, 604, L73
- Mao, S. & Schneider, P. 1998, MNRAS, 295, 587
- Mao, S., Jing, Y., Ostriker, J. P., & Weller J., 2004, ApJ, 604, L5
- McGaugh, S. S. & de Blok, W. J. G. 1998, ApJ, 499, 41

- Metcalf, R. B. & Madau, P. 2001, ApJ, 563, 9
- Metcalf, R. B. & Zhao, H. 2002, ApJ, 567, L5
- Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J., & Tozzi, P. 1999, ApJ, 524, L19
- Natarajan, P. & Kneib, J. 1997, MNRAS, 287, 833
- Natarajan, P., Kneib, J., Smail, I., & Ellis, R., 2004, submitted
- Natarajan, P., Kneib, J., Smail, I., & Ellis, R. S. 1998, ApJ, 499, 600
- Natarajan, P., Loeb, A., Kneib, J., & Smail, I. 2002, ApJ, 580, L17
- Navarro, J. F., Hayashi, E., Power, C., Jenkins, A. R., Frenk, C. S., White, S. D. M., Springel, V., Stadel, J., & Quinn, T. R. 2004, MNRAS, 349, 1039
- Springel, V., White, S. D. M., Tormen, G., & Kauffmann, G. 2001a, MNRAS, 328, 726
- Springel, V., Yoshida, N., & White, S. D. M. 2001b, New Astronomy, 6, 79
- Stoehr, F., White, S. D. M., Tormen, G., & Springel, V. 2002, MNRAS, 335, L84
- Taylor, J. E. & Babul, A. 2001, ApJ, 559, 716
- Trentham, N., Möller, O., & Ramirez-Ruiz, E. 2001, MNRAS, 322, 658
- Wilson, G., Kaiser, N., & Luppino, G. A. 2001, ApJ, 556, 601
- Zentner, A. R. & Bullock, J. S. 2003, ApJ, 598, 49




Fig. 1.— The substructure mass function retrieved from galaxy-galaxy lensing for 5 massive HST cluster-lenses. This mass spectrum is constituted by the dark halos associated with the 40 brightest galaxies in these clusters. The panel on the bottom right shows the coadded and averaged clump spectrum of clusters A2218, A2390, and Cl0054.




Fig. 2.— Comparison of the substructure mass function of the stacked clusters A2210, A2390, and Cl0054 (shaded histogram) with N-body simulations of a rich cluster of galaxies (filled histogram), carried out at four different numerical resolutions.